

École Doctorale des Sciences de l'Environnement d'Île-de-France

Année Universitaire 2021-2022

Modélisation Numérique
de l'Écoulement Atmosphérique
et Assimilation de Données


Olivier Talagrand


Cours 1

17 Mars 2022

Programme of the course

1. Numerical modelling of the atmospheric flow. The *primitive* equations. Discretization methods. Numerical Weather Prediction. Present performance.
2. The meteorological observation system. The problem of 'assimilation'. Bayesian estimation. Random variables and random functions. Meteorological examples.
3. 'Optimal Interpolation'. Basic properties. Meteorological applications. The theory of *Best Linear Unbiased Estimator*.
4. Advanced assimilation methods.
 - Kalman Filter. Ensemble Kalman Filter. Present performance and perspectives.
 - Variational Assimilation. Adjoint Equations. Present performance and perspectives.
5. Advanced assimilation methods (continuation).
 - Bayesian Filters. Theory, present performance and perspectives.





Bilan radiatif de la Terre, moyenné sur un an

Global Energy Flows $W m^{-2}$






FIG. 1. The global annual mean Earth's energy budget for the Mar 2000 to May 2004 period ($W m^{-2}$). The broad arrows indicate the schematic flow of energy in proportion to their importance.



Annual mean energy budget of atmosphere for period 2000–2010. Unit $W \cdot m^{-2}$

Stephens *et al.*, *Nature Geoscience*, 2012





Particle moves on sphere with radius R
under the action of a force lying
in meridian plane of the particle

→ Angular momentum wrt axis of rotation conserved.

$$(u + \Omega R \cos\varphi) R \cos\varphi = Cst$$

On Earth, $\Omega \approx 2\pi \cdot 10^{-5} \text{ s}^{-1}$, $R \approx 6.4 \cdot 10^6 \text{ m}$.

If $u = 0$ at equator, $u = 329 \text{ ms}^{-1}$ at latitude $\varphi = 45^\circ$. If $u = 0$ at 45° , $u = -232 \text{ ms}^{-1}$ at equator.

Hadley, G., 1735, Concerning the cause of the general trade winds, *Philosophical Transactions of the Royal Society*

The general circulation




Fig. 10.1. Streak photographs illustrating the dependence of the flow type on rotation rate Ω for a laboratory 'dishpan' experiment. The values of Ω in rad s^{-1} are (a) 0.41; (b) 1.07; (c) 1.21; (d) 3.22; (e) 3.91; (f) 6.4. Working fluid was a water-glycerol solution of mean density 1.037 g cm^{-3} and kinematic viscosity $1.56 \times 10^{-2} \text{ cm}^2 \text{ s}^{-1}$. The streak photographs show the flow at a depth of 0.5 cm below the free upper surface (see also problem 10.1.) (From Hide & Mason, 1975)




Figure 1.3. Schematic regime diagram for the thermally driven rotating annulus in relation to the thermal Rossby number Θ (or stability parameter, $\propto \Omega^{-2}$) and Taylor number $\mathcal{T} \propto \Omega^2$, showing some typical horizontal flow patterns at the top surface, visualized as streak images at upper levels of the experiment.

Read *et al.*, 2015





Figure 2. Comparison shows similarities between the global 500 mb pressure pattern in the upper atmosphere of the Northern Hemisphere and a four-wave pattern in the laboratory.


(Laboratory flow conditions were similar to those in Fig. 1, except $\Omega = 1.95$ radians per sec.) In the atmosphere the flow is approximately parallel to the isobars (the flow is to the right,



from high to low pressure), with speed inversely proportional to the spacing. Changes in the wave pattern have a significant effect on large-scale weather and climate.




26/04/1984, 00/00 TU



Zzonal wind; annual longitudinal average (m.s^{-1})



<http://paoc.mit.edu/labweb/notes/chap5.pdf>,

Atmosphere, Ocean and Climate Dynamics, by J. Marshall and R. A. Plumb,
International Geophysics, Elsevier)




Peixoto and Oort, 1992, *The Physics of Climate*, Springer-Verlag



Base time: Fri 11 Mar 2022 00 UTC Valid time: Wed 16 Mar 2022 00 UTC (+120h) Area : Europe



Base time: Wed 16 Mar 2022 00 UTC Valid time: Wed 16 Mar 2022 00 UTC (+0h) Area : Europe



Base time: Fri 11 Mar 2022 12 UTC Valid time: Fri 11 Mar 2022 12 UTC (+0h) Area : Europe



Base time: Wed 16 Mar 2022 00 UTC Valid time: Wed 16 Mar 2022 00 UTC (+0h) Area : Europe






Fig. 1: Members of day 7 forecast of 500 hPa geopotential height for the ensemble originated from 25 January 1993.



Figure 6 Hurricane Katrina mean-sea-level-pressure (MSLP) analysis for 12 UTC of 29 August 2005 and $t+84h$ high-resolution and EPS forecasts started at 00 UTC of 26 August:


- 1st row: 1st panel: MSLP analysis for 12 UTC of 29 Aug
 2nd panel: MSLP $t+84h$ T₁₅₁L60 forecast started at 00 UTC of 26 Aug
 3rd panel: MSLP $t+84h$ EPS-control T₂₅₅L40 forecast started at 00 UTC of 26 Aug
 Other rows: 50 EPS-perturbed T₁₂₅₅L40 forecast started at 00 UTC of 26 Aug.

The contour interval is 5 hPa, with shading patterns for MSLP values lower than 990 hPa.

Ensemble forecast at range 7 days

ECMWF ENSEMBLE FORECASTS

Sunday 06 March 2022 1200 UTC ECMWF forecast 1-168 VT: Sunday 13 March 2022 1200 UTC
MSLP (contour every 5hPa) Temperature at 850hPa (only -4 and 16 isolines are plotted)



Pourquoi les météorologistes ont-ils tant de peine à prédire le temps avec quelque certitude ? Pourquoi les chutes de pluie, les tempêtes elles-mêmes nous semblent-elles arriver au hasard, de sorte que bien des gens trouvent tout naturel de prier pour avoir la pluie ou le beau temps, alors qu'ils jugeraient ridicule de demander une éclipse par une prière ? Nous voyons que les grandes perturbations se produisent généralement dans les régions où l'atmosphère est en équilibre instable. Les météorologistes voient bien que cet équilibre est instable, qu'un cyclone va naître quelque part ; mais où, ils sont hors d'état de le dire ; un dixième de degré en plus ou en moins en un point quelconque, le cyclone éclate ici et non pas là, et il étend ses ravages sur des contrées qu'il aurait épargnées. Si on avait connu ce dixième de degré, on aurait pu le savoir d'avance, mais les observations n'étaient ni assez serrées, ni assez précises, et c'est pour cela que tout semble dû à l'intervention du hasard.

H. Poincaré, *Science et Méthode*, Paris, 1908

Why have meteorologists such difficulty in predicting the weather with any certainty? Why is it that showers and even storms seem to come by chance, so that many people think it quite natural to pray for rain or fine weather, though they would consider it ridiculous to ask for an eclipse by prayer? We see that great disturbances are generally produced in regions where the atmosphere is in unstable equilibrium. The meteorologists see very well that the equilibrium is unstable, that a cyclone will be formed somewhere, but exactly where they are not in a position to say; a tenth of a degree more or less at any given point, and the cyclone will burst here and not there, and extend its ravages over districts it would otherwise have spared. If they had been aware of this tenth of a degree they could have known it beforehand, but the observations were neither sufficiently comprehensive nor sufficiently precise, and that is the reason why it all seems due to the intervention of chance.

H. Poincaré, *Science et Méthode*, Paris, 1908
(English transl. by F. Maitland, *Science and Method*,
T. Nelson and Sons, London, 1914)

Physical laws governing the flow

- Conservation of mass

$$D\rho/Dt + \rho \operatorname{div}\underline{U} = 0$$

- Conservation of energy

$$De/Dt - (p/\rho^2) D\rho/Dt = Q$$

- Conservation of momentum

$$D\underline{U}/Dt + (1/\rho) \operatorname{grad}p - \underline{g} + 2 \underline{\Omega} \wedge \underline{U} = \underline{F}$$

- Equation of state

$$f(p, \rho, e) = 0 \quad (p/\rho = rT, e = C_v T)$$

- Conservation of mass of secondary components (water, chemical species, ...)

$$Dq/Dt + q \operatorname{div}\underline{U} = S$$

These physical laws must be expressed in practice in discretized (and necessarily imperfect) form, both in space and time

$$Da/Dt = \partial a/\partial t + \underline{U \cdot \text{grad} a}$$

Particular
(Lagrangian)
derivative

Eulerian
derivative

Advection
(due to motion)

The case of the ocean

- Same basic equations

But

- Different equation of state
- Major secondary component is now salt (convective instability associated with variation of density of the fluid)

Physical laws must in practice be discretized in both space and time
 \Rightarrow *numerical models*, which are necessarily imperfect.

Models that are used for large scale weather prediction and for climatological simulation cover the whole volume of the atmosphere. These models are based, at least so far, on the *hydrostatic* hypothesis

in the vertical direction :

$$\partial p / \partial z + \rho g = 0$$

Eliminates momentum equation for vertical direction. In addition, flow is incompressible in coordinates (x, y, p) \Rightarrow number of equations decreased by two units.

Hydrostatic approximation valid, to accuracy $\approx 10^{-4}$, for horizontal scales
> 20-30 km

More costly nonhydrostatic models are used for small scale meteorology.

Hydrostatic approximation allows to take pressure p as independent vertical coordinate

- Flow is incompressible

- Pressure gradient term $(1/\rho) \text{grad}_z p$ becomes $\text{grad}_p \Phi$, where $\Phi \equiv gz$ is geopotential


Spatial Discretization

There exist at present two forms of spatial discretization

- Gridpoint discretization
- (Semi-)spectral discretization (mostly for global models, and most often only in the horizontal direction)


Finite element discretization, which is very common in many forms of numerical modelling, is rarely used for modelling of the atmosphere, except for discretization in the vertical direction. It is more frequently used for oceanic modelling, where it allows to take into account the complicated geometry of coast-lines.

In gridpoint models, meteorological fields are defined by values at the nodes of a grid covering the physical domain under consideration. Spatial derivatives are expressed by finite differences.




$$\left(\frac{\partial \Phi}{\partial x} \right)_{i,j} \approx \frac{\Phi_{i+1,j} - \Phi_{i-1,j}}{2\Delta x}$$


A schematic of an Atmospheric General Circulation Model (L. Fairhead /LMD-CNRS)



Grille Emerald-Péridot



Grille Arpège



Grilles de modèles de Météo-France (*La Météorologie*)

In *spectral models*, fields are defined by the coefficients of their expansion along a prescribed set of basic functions. This is similar to what is often done in a periodic domain in R^n , by taking imaginary exponential functions (sines and cosines) as basis functions

$$F(x, y) = \sum_{k, m} \mathcal{F}(k, m) \exp [2i\pi (kx/L_x + my/L_y)]$$

where the function F has periods L_x and L_y in the directions x and y respectively.

In the discretized case, the integer indices k and l are limited to a set of finite values

$$k = -K, \dots, 0, \dots, K \quad ; \quad m = -M, \dots, 0, \dots, M$$

Advantage : spatial differentiation is obvious and exact

In the case of global meteorological models, which cover the whole atmosphere, the basis functions are the *spherical harmonics*

$$T(\mu=\sin(\text{latitude}), \lambda=\text{longitude}) = \sum_{\substack{0 \leq n < \infty \\ -n \leq m \leq n}} T_n^m Y_n^m(\mu, \lambda)$$

où les $Y_n^m(\mu, \lambda)$ sont les *harmoniques sphériques*

$$Y_n^m(\mu, \lambda) \propto P_n^m(\mu) \exp(im\lambda)$$

$P_n^m(\mu)$ est la *fonction de Legendre* de deuxième espèce


$$P_n^m(\mu) \propto (1 - \mu^2)^{\frac{m}{2}} \frac{d^{n+m}}{d\mu^{n+m}} (\mu^2 - 1)^n$$

n et m sont respectivement le *degré* et l'*ordre* de l'harmonique $Y_n^m(\mu, \lambda)$


$$n = 0, 1, \dots \quad -n \leq m \leq n$$

Годн и изобразя, ие две сферически


$$l = 3$$
$$m = 0$$
$$l - m = 3$$




$$l = 3$$
$$m = 1$$
$$l - m = 2$$



$$l = 3$$
$$m = 2$$
$$l - m = 1$$



$$l = 3$$
$$m = 3$$
$$l - m = 0$$



$$l = 5$$
$$m = 2$$
$$l - m = 3$$



Les harmoniques sphériques sont fonctions propres du laplacien à la surface de la sphère

$$\Delta Y_n^m = -n(n+1)Y_n^m$$

Troncature 'triangulaire' TN ($n \leq N, -n \leq m \leq n$) indépendante du choix d'un axe polaire. Représentation est parfaitement homogène à la surface de la sphère

Calculs non linéaires effectués dans l'espace physique (sur grille appropriée, souvent latitude-longitude 'gaussienne'; nécessaire pour éviter stroboscopie, ou *aliasing*). Les transformations requises sont possibles à un coût non prohibitif grâce à l'utilisation de Transformées de Fourier Rapides (*Fast Fourier Transforms, FFT*, en anglais). Il existe aussi une version rapide des Transformées de Legendre, relatives à la variable μ .

Du fait de ces transformations permanentes entre l'espace spectral et l'espace physique, ces modèles sont souvent appelés *semi-spectraux*.

In addition to hydrostatic approximation, the following approximations are (almost) systematically made in global modeling :

- Atmospheric fluid is contained in a spherical shell with negligible thickness. This does not forbid the existence within the shell of a vertical coordinate which, in view of the hydrostatic equation, can be chosen as the pressure p .

- The horizontal component of the Coriolis acceleration due to the vertical motion is neglected (this approximation, sometimes called the *traditional approximation*, is actually a consequence of the previous one).

- Tidal forces are neglected.

These approximations lead to the so-called (and ill-named) *primitive equations*

Pressure p , although convenient for writing down the equations, is in fact rather inconvenient because lower boundary is not fixed in (x, y, p) -space.

So-called σ -coordinate. $\sigma \equiv p/p_S$, where p_S is pressure at ground level.

‘Hybrid’ coordinate.

Physical laws governing the flow

- Conservation of mass

$$D\rho/Dt + \rho \operatorname{div}\underline{U} = 0$$

- Conservation of energy

$$De/Dt - (p/\rho^2) D\rho/Dt = Q$$

- Conservation of momentum

$$D\underline{U}/Dt + (1/\rho) \operatorname{grad}p - \underline{g} + 2 \underline{\Omega} \wedge \underline{U} = \underline{F}$$

- Equation of state

$$f(p, \rho, e) = 0 \qquad (p/\rho = rT, e = C_v T)$$

- Conservation of mass of secondary components (water in the atmosphere, salt in the ocean, chemical species, ...)

$$Dq/Dt + q \operatorname{div}\underline{U} = S$$

These physical laws must be expressed in practice in discretized (and necessarily imperfect) form, both in space and time


Parlance of the trade :

One ordinarily distinguishes two different parts in models. The ‘**dynamics**’ deals with the physically reversible processes (pressure forces, Coriolis force, advection, ...), while the ‘**physics**’ deals with physically irreversible processes, in particular the diabatic heating term Q in the energy equation, and also the parameterization of subgrid scales effects.

Numerical schemes have been gradually developed and validated for the ‘dynamics’ component of models, which are by and large considered now to work satisfactorily (although regular improvements are still being made; project *DYNAMICO*, *Dynamical Core on Icosahedral Grid*, Th. Dubos, IPSL).

The situation is different as concerns 'physics', where many problems remain (as concerns for instance subgrid scales parameterization, the water cycle and the associated exchanges of energy, or the exchanges that take place in the boundary layer between the atmosphere and the underlying medium). 'Physics' as a whole remains the weaker point of models, and is still the object of active research.

5 - SCHEMA DES INTERACTIONS PHYSIQUES DANS LE MODELE



Temporal Discretization

Equation

$$dx / dt = F(x)$$

(x state vector of the model).

Timestep Δt .

Computed solution at time $n\Delta t$ denoted x_n

Forward (Euler) scheme

$$(x_{n+1} - x_n)/\Delta t = F(x_n)$$

$$x_{n+1} = x_n + \Delta t F(x_n)$$

Implemented on equation

$$dx/dt = i\alpha x \quad , \quad \alpha \text{ real} \quad (1)$$

Exact solution $x(t) = x(0) \exp(i\alpha t)$

Modulus $|x(t)|$ conserved in time

Discretized solution according to forward scheme

$$x_{n+1} = (1 + i\alpha\Delta t) x_n$$

Modulus $|x_{n+1}| = \sqrt{1 + \alpha^2\Delta t^2} |x_n|$

increases exponentially with time.

Forward scheme is *unconditionally unstable* for Eq. (1)

Leapfrog scheme

$$(x_{n+1} - x_{n-1})/2\Delta t = F(x_n)$$

$$x_{n+1} = x_{n-1} + 2\Delta t F(x_n)$$

Stable for equation (1) above (*i.e.* modulus remains constant in time) provided

$$\alpha\Delta t < 1$$

Courant-Friedrichs-Lewy (CFL) condition

In a multidimensional system, the largest α will be the highest frequency that is present in the system. In a discretized system of travelling waves, the highest frequency will correspond to the fastest wave that the discretization can explicitly resolve. It will be proportional to $c/\Delta x$, where c is the phase velocity of the fastest waves in the system, and Δx the mesh-size of the discretization

$$\alpha = (1/\beta) c/\Delta x$$

where β is an $O(1)$ numerical coefficient depending on the particular discretization scheme under consideration.

CFL condition then becomes

$$\Delta t / \Delta x < \beta / c$$

Significance : numerical propagation of signal must be at least as fast as physical propagation.

CFL condition generally applies to explicit schemes of temporal discretization

In hydrostatic atmosphere, fastest propagating wave : gravity wave with largest scale height, $c = \sqrt{rT} \approx 300 \text{ m.s}^{-1}$.

$$\Delta x = 30 \text{ km} \quad \Rightarrow \quad \Delta t = 100 \text{ s}$$

The use of *semi-implicit* schemes allows to get rid of the CFL condition, and to use longer timesteps.

Cours à venir

~~Jeudi 17 mars~~

Jeudi 24 mars

Jeudi 31 mars

Jeudi 14 avril

Jeudi 21 avril

Jeudi 28 avril

Jeudi 5 mai

Jeudi 12 mai