

A new look at the atmospheric water cycle: measurements of water vapor and its main isotopologue using SCIAMACHY

Christian Frankenberg (1), Remco Scheepmaker (2), Ilse Aben (2), Hans Schrijver (2), Annemieke Gloudemans (2), Thomas Roeckmann (3), and Kei Yoshimura (4)

(1) Jet Propulsion Laboratory, California Institute of Technology, Pasadena, USA

(2) SRON Netherlands Institute for Space Research, Utrecht, The Netherlands

(3) Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, The Netherlands

(4) Scripps Institution of Oceanography, University of California, La Jolla, USA

Water vapor is by far the most important greenhouse gas in the atmosphere. As a warmer atmosphere can contain more water vapor, a positive feedback effect with respect to climate change is expected. The distribution of water vapor is very inhomogeneous and variable, unlike that of other greenhouse gases. In the light of climate reconstructions and predictions, it is therefore crucial to better understand the water cycle and its response to past and present climate change. The relative abundance of the heavy water isotopologue HDO provides a deeper insight in the water cycle, as evaporation and condensation processes deplete heavy water in the gas phase. In the application of isotopologues, however, the space-borne retrieval of atmospheric water vapor isotopologues near the surface has so far been overlooked. We provide, for the first time (Frankenberg et al., Science 2009), global HDO/H₂O abundances using the Scanning Imaging Absorption spectrometer for Atmospheric CHartography (SCIAMACHY) instrument onboard ENVISAT. This allows for an entirely new perspective on the near-surface distribution of water vapor isotopologues. We are using the 2.3 micron (SWIR) window of SCIAMACHY, which is also used for the first time (Schrijver et al., AMT 2009) to derive total water vapor columns. Using absorption spectroscopy in the short-wave infrared, we are sensitive down to the lowest parts of the atmosphere where most of the water vapor resides. We will present the results shown in the recent publication, retrieval methods, potential applications in hydrological and atmospheric research as well as future potentials.